

The “Efucal” combination

Efucal contains two main ingredients:

1. **5-fluorouracil:** Also known as 5-FU or *Efudix*, this drug has been used for many years as an effective treatment for solar keratoses and Bowen’s disease. It is toxic to cancerous and pre-cancerous cells, but has little effect on normal undamaged skin.
2. **Calcipotriol :** This is a form of vitamin D which stimulates the immune system to attack cancer and pre-cancer cells.

In combination, this treatment has significant advantages over 5-fluorouracil alone:

- The treatment course lasts days rather than weeks
- There is less inflammation and irritation than when 5-fluorouracil is used alone
- The treated area is less likely to develop squamous cell carcinoma for at least 3 years after treatment (with 5-fluorouracil alone, this benefit lasts only one year).

Where are some places you can get Efucal

CHEMPRO (07) 3849 6938
1173 Logan Rd, Holland Park QLD 4121

CHEMPRO– (07) 3343 9330
MOUNT GRAVATT PLAZA

East Coast Compounding (07) 3822 6444
687-689 Old Cleveland Rd East, Wellington Point 4160

How to take Efucal

Solar Keratosis treatment length with Efucal

Note that the duration of treatment many need to be varied depending on clinical response or side effects.

Face, chest and scalp	Twice daily for 4 days (6-8 days if first attempt too weak)
Lip	Twice daily for 1-2 days
Other areas, e.g. hands, forearms, legs	Twice daily for 6 days (8-14 days if first attempt too weak)
OFF LABEL treatment of Bowen's disease (early SCC skin cancer)	Twice daily for 4 days. 14 day break. Twice daily for a further 4 days. Note: Further treatment may be required depending on treatment response.

Wait 8 weeks before deciding on whether need to retreat.

Application instructions

1. Wash the skin to be treated using lukewarm water. Avoid soap which may sting or irritate your skin, especially as treatment progresses.
2. Pat dry the affected area.
3. Allow your skin to completely dry.
4. Apply a thin layer of cream to the affected area(s) twice a day, without a dressing, for the duration advised above. Do not continue further unless advised.
5. Use a non-metal applicator, cotton bud or a rubber glove to apply a thin film of cream to your affected skin. Use the smallest amount that can cover the treatment area.
6. Don't allow the cream to contact mucous membranes, such as eyes, eyelids, nostrils and lips, unless we are specifically targeting these areas.
7. Do not apply to broken skin or open cuts as it may be absorbed into the bloodstream and may cause side effects.
8. Wash your hands carefully and thoroughly immediately after using the cream.
9. Do not cover the treated area unless advised.

Apply sunscreen 20 minute after morning application

Apply night-time dose at least 2 hrs before going to bed

The total area of skin treated with at any time should not exceed 500 cm² (approx. 23 x 23cm). Treat larger areas a section at a time.

Provoking stronger response (OPTIONAL)

For thick scaly solar keratosis, previously treated solar keratosis with a poor response or when treating the arms or legs you may need to prime the skin with a [strong steroid](#) like clobetasol/diprosone. If pre-treating with steroid, apply the steroid to the skin twice daily for 10 days before treatment.

Can also use an extra 10-15% calcipotriol to boost the effect of the efucal if stronger treatment response desired.

Precautions

Efucal must not be used during pregnancy or while breastfeeding.

Where possible avoid sun exposure to the treated area both during treatment and while the skin is irritated and inflamed after treatment. Even untreated areas of the skin may be more sensitive to sunlight.

What to expect

Usual Course of events:

Redness day 4-5

Stinging and itch day 6

Resolution day 14-16

Occasional severe treatment effects occur – even after the first application which may lead to prolonged crusting, cracked skin and bleeding

This means a stronger response but is more painful and longer lasting

The treated area while appear unsightly and may be uncomfortable during treatment and for up to 2 weeks after the course is completed.

Side effects and aftercare

Irritated skin may be itchy or painful. There may be a burning sensation, redness and ulcers or sores in the treated area. Sometimes these symptoms affect areas that did not previously have solar keratoses. This means that the ingredients are treating sun damaged skin that has not yet developed into solar keratoses or skin cancer.

Paraffin ointment/petroleum jelly (e.g. [Bepanthen](#), [pawpaw ointment](#) or [Vaseline](#)) can provide relief. For further relief store these treatments in the fridge.

For severe treatment effects e.g. pain responding to above treatments, concerning skin inflammation, cracked and bleeding skin OR a need to speed resolution of treatment effects a [strong steroid ointment](#) like clobetasol can be used twice daily to the affected areas for 2-4 days. Severe treatment effect suggests the treatment has been extremely effective. Using a strong steroid will dampen this somewhat – however the treatment will still be effective and is encouraged if the side effects are too severe, especially if you are worried about scarring of the skin (rare).

Do not use antiseptics such as Betadine, as they tend to dry the skin and make irritation and pain worse. If you suspect that the skin has become infected, notify your doctor.

In unusual cases, more general side effects can occur. These include fever, dizziness, abdominal pain and cramps, diarrhoea and vomiting. If you experience any of these symptoms, stop using Efucal and notify your doctor.

Storage

Store at room temperature